

PERIODONTAL DISEASES AND ORAL HYGIENE PRACTICES AMONG A GROUP OF DRUG ADDICTED PRISONERS

Shailendra Singh Chaturvedi,¹ Hiroj Bagde²
Reader,¹ Senior Lecturer²

1 – 2 Department of Periodontics, New Horizon Dental College & Research Institute, Bilaspur.

Abstract

Purpose: The present study aimed to assess the relation of periodontitis with respect to different factors among this population.

Methods: The study sample consisting of 200 drug addict male subjects with mean age of 41.32 ± 2.32 years. A self-administered structured questionnaire validated through a pilot survey, using WHO Oral Health Assessment Form (1997) was prepared. Periodontal status was assessed by using CPI probe. Chi-square and ANOVA test were used to find the significance of difference at p value <0.05 .

Results: The information gathered regarding oral hygiene practices showed that most of the subjects (36.60%) used finger and powder to maintain their oral hygiene. The overall prevalence of periodontal diseases was high with common occurrence of shallow and deep pockets. Karl Pearson's correlation also showed a linear relationship of age with periodontal diseases. (CPI & LOA)

Conclusion: There is an attentive need of oral health care programmes as the standard of periodontal health observed among inmates is low.

Key words: Adverse Habits, Inmates, Periodontitis, Treatment Needs.

Introduction

Despite great achievements in the oral health of populations globally, problems still remain in many communities around the world, particularly among the disadvantaged and socially marginalized groups (WHO oral Health Report 2003).¹ Prisoners constitute one of the disadvantaged group as these people are solitary confined for long duration and hence are socially and economically deprived from the community. Prison population is unique and challenging with many oral health problems. Dental diseases can reach epidemic proportions in the prison setting.² The inmate population in India has increased faster over the years as there has been shift towards a more penalizing response to crime, resulting in longer sentences. The National Crime Bureau Report in 2010 suggested that the number of prisoners in the country is estimated to be 3,68,998 whereas capacity in jails is 3,20,450.³

Hence the overcrowding, violence and isolation may have an impact on both general as well as oral health status of inmates.⁴ Furthermore most of them had been already involved in adverse habits like drug addiction and develop mental health problems due to its withdrawal symptoms. Drug abuse results in various individual and social consequences and takes a heavy toll in terms of severe health complications.⁵ It also modifies the behaviour and lifestyle of the person. Moreover factors like nutritional status, tobacco consumption, poor oral hygiene, stress, negligible dental visits are linked to a wide range of periodontal diseases forming the fundamental basis of the common risk factor.⁶ Psychological factors are also considered as the most common cause of development of para-functional habits like bruxism, nail biting, tongue thrusting etc. which in turn can affect the periodontium. Hence inmates are more vulnerable to periodontal diseases due to the above mentioned factors.⁷ Various studies have been done on the oral health conditions of the general

population.^{8,9} But limited data published regarding oral health status of prisoners,¹⁰ may be due to lack of dental staff in the prison setting. Therefore, this study has been done to assess the prevalence of periodontal diseases among this population.

Methodology

The total study population of the jail was around 970 and around 230 were in the de-addiction centre. Those who agreed to participate in the survey and had been involved with habits like alcoholism, tobacco consumption, drug addiction were taken in the study. Prisoners who were uncooperative and were suffering from other systemic disorders like cardiovascular problems were excluded.

Hence the study comprised of 200 male inmates aged above 25 years with a mean age was 41.32 ± 2.32 years. Official permission was obtained from the Head of the Institute and jail authorities before conducting the study and a written informed consent was obtained from all the participants. A self-administered structured questionnaire validated through a pilot survey, consisting of 3 sections: Information regarding general demographic variables; Questions regarding oral hygiene practices; and Assessment of the periodontal status using WHO Oral Health Assessment Form (1997).¹¹ The oral examination was done using standard explorers, mouth mirrors and CPI probes.

The intra-examiner reliability for various recordings ranged from 0.90 to 0.96. Proper sterilization was maintained during the examination.


Data analysis

The Statistical software namely SPSS version 16.0 was used for the analysis of the data. Chi-square and ANOVA test were used to find the significance of difference among different age groups at p value <0.05 . Karl Pearson's

correlation was used to assess the relationship of age with periodontal diseases.


Results

The information gathered regarding oral hygiene practices among the participants showed that 30.30% subjects used tooth brush with tooth paste, 36.60% used finger and powder, and 28.70% used chew sticks whereas 4.40% infrequently clean their teeth. (Graph 1)



Graph 1: Distribution of study subjects according to oral hygiene practices

Similarly, a significant difference with highest Loss of attachment was observed in different age groups. Maximum number of subjects between the age group of 25-34 years (77.10%) had no loss of attachment. Loss of attachment of 4-5 mm was seen more among 35-44 year age group i.e. 19.40% whereas loss of attachment of 6-8mm, 9-11mm and 12mm or more were found more in the oldest age group (≥ 45 years) i.e. (25.00%), (14.30%) and (3.60%) respectively. (Graph 2)



Graph 2: Showing study population with highest percentage of Loss of Attachment according to age

The overall mean number of sextants for bleeding component was 1.68 1.549, that of calculus 1.59 1.625, for shallow pockets (4-5 mm) 0.92 1.156 and for deep

pockets (6mm or more) 0.30 0.682. The distribution of mean number of sextants according to age groups showed that as the age advances bleeding scores declines whereas calculus and pocket formation increases significantly. (Table 1)

| Age groups | No | Bleeding | | Calculus | | Pocket (4-5mm) | | Pocket (6mm or more) | |
|-----------------|-----|----------|-------|----------|-------|----------------|-------|----------------------|-------|
| | | Mean | SD | Mean | SD | Mean | SD | Mean | SD |
| 25-34 years | 76 | 2.15 | 1.774 | 1.08 | 1.318 | .40 | .610 | .08 | .279 |
| 35-44 years | 65 | 1.42 | 1.339 | 2.06 | 1.866 | .72 | .974 | .33 | .632 |
| ≥ 45 years | 59 | 1.21 | 1.166 | 1.86 | 1.580 | 2.07 | 1.303 | .64 | 1.026 |
| Total | 200 | 1.68 | 1.549 | 1.59 | 1.625 | .92 | 1.156 | .30 | .682 |
| F value | | 4.182 | | 4.449 | | 29.191 | | 6.601 | |
| P value | | 0.018* | | 0.014* | | 0.000* | | 0.002* | |

Table 1: Mean number of sextants affected by periodontal disease according to age

| | Age | Code CPI | Code LOA |
|---------------------|-----|----------|----------|
| Pearson Correlation | 1 | .400(*) | .321(*) |
| Sig. (2-tailed) | | .000 | .000 |

Table 2 shows a linear relationship of age with Community Periodontal Index (CPI) and Loss of Attachment (LOA) using Karl Pearson's correlation coefficient ($p = 0.000$).

Discussion

The current study population is mostly from the lower socioeconomic group which is deprived from the society. So their isolated environment makes them more depressed which affect the systemic as well as oral health of the body and problems like periodontitis become more prevalent.(12) This study is the first to present information about periodontal health among drug addicted inmates. The majority of the participants belonged to age group of 25-34 years with overall mean age of all the subjects was 39.46 ± 2.28 years and the observations were similar to other studies done by Reddy *et al.* (2012), Brijender *et al.* (2006) and Jagadeesan *et al.* in 2003.¹³ Almost all the participants in the present study had been involved in the adverse habits such as substance use and tobacco consumption. However the prevalence of drug or alcohol consumption found to be approximately 70% among inmates in Canada.¹⁴ Heidari *et al* found its usage in 83% of remand prisoners in London.¹⁵ The higher prevalence of adverse habits in the present data could be due to the selected sample as it was taken from the de-addiction centre from the jail. Other reason could be of unwillingness to report about drug addiction when surveyed. According to oral hygiene behaviour, most of the prisoners used finger & powder (36.6%) which is comparable to Punitha *et al* study in 2011 among rural

children. The similarity could be because of lower level of knowledge in young age especially in rural regions.⁸ The percentage of chew stick users was 28.7% which was much lower than Bhat *et al* study among tribal community.⁹ It could be due to less availability of sticks in the jail premises. And twigs of the arak tree (*S.persica*) had been used for centuries as oral hygiene tools. So the tribal population is still having the same attitude towards oral health.¹⁶ But it was also observed that the participants do oral hygiene practices only for namesake as most of them are illiterate and show little concern for cleanliness. The jail inmates showed pocket formation of 4-5 mm and 6mm or more as 0.92 ± 1.156 and 0.30 ± 0.682 respectively. Whereas Singh *et al* found higher scores of mean sextants among prisoners at Lucknow jail in 2012. It could be due to more prevalence of adverse habits among people residing in Lucknow area than the present study locality.¹⁰ However other study conducted by Jain *et al* (2009) has shown higher score for 4-5 mm pockets and lower for 6 mm or more pockets.¹⁷ Similarly many studies have shown lower scores than the present findings^{18,19,20} which may be due to many factors as the inmates had experienced many adverse substances. And the withdrawal symptoms increase the stress level that affects periodontal health. Regarding loss of attachment, 19.40% of the population in the present study suffered from loss of attachment of 4-5 mm followed by 13.90% having 6-8mm and 5.60% having 9-11mm. These findings are more than those conducted by Bali RK *et al* (2003) among Indian population of similar age group (35 to 44) years, could be because of poor quality of life among this community.²¹ Overall loss of attachment in this study was more than Singh SK *et al* findings.¹⁰ It could be due to habit of using chew sticks among the subjects that leads to recession.

According to Reddy *et al* 66.3% of inmates had poor oral hygiene status; 39.3% had a CPI score of 2 and 48.6% had a CPI score of 3 or 4; 30.1% of inmates had a loss of attachment (LOA) score of 1 or 2 (more than 4 mm) and 1.7% of inmates had a score of 4.²² Moreover, high correlation has been documented between age and the progression of periodontal disease, in which the greatest pocket depth scores were in the older prisoners.¹⁵ This possibility may be due to ignorance of dental care as people think that any how teeth will become loose with advancing age.⁹ Additionally, all the negative factors of health lead to malnutrition and making them more susceptible for infections. Moreover, a high percentage of the study population accounted dependency on drugs, and this may impact on oral health as a result of the high levels of dental anxiety, and avoidance of dental treatment reported amongst drug users.

This susceptibility coupled with poor oral hygiene, leads to a higher prevalence of periodontal diseases which has been well documented in other studies also.^{22,23,24,25}

Conclusion

The results of the present study emphasized a need for special attention from government sector and other health

organizations to meet the basic oral health needs of this community as high prevalence of periodontitis was observed which ultimately showed positive correlation with lack of oral hygiene practices, deprivation of social life, and poor nutrition and overall quality of life in the jail setting. Also oral health care programmes should be organized for prisoners in order to educate them regarding proper oral hygiene practices.

References

1. Petersen PE. Improvement of oral health in Africa in the 21st century – the role of the WHO Global Oral Health Programme. *Developing Dentistry*. 2004;5(1):9-20.
2. Freudenberg N. Jails, Prisons and the Health of Urban Populations: A Review of the Impact of the Correctional System on Community Health. *J Urban Health* 2001;78(2):214-234.
3. Das SK. Report of the Committee on Crime Statistics. Social Statistics Division Central Statistics Office Ministry of Statistics & Programme Implementation. Sardar Patel Bhavan, Sansad Marg New Delhi – 110001.
4. Levy MH. Australian prisons are still health risks. *Med J Aust* 1999;171(1):7-8.
5. Jernigan DH, Monteiro M, Room R, Saxena S. Towards a global alcohol policy: alcohol, public health and the role of WHO. *Bull World Health Organ* 2000;78(4):491-499.
6. Gjermo P, Rosing CK, Susin C, Oppermann R. Periodontal diseases in Central and South America. *Periodontol* 2000. 2002;29:70–78.
7. Morio KA, Marshall TA, Qian F, Morgan TA. Comparing diet, oral hygiene and caries status of adult methamphetamine users and nonusers: a pilot study. *J Am Dent Assoc* 2008;139(2):171-176.
8. Punitha VC, Sivaprakasam P. Oral Hygiene Status, Knowledge, Attitude and Practices of Oral Health among Rural Children of Kanchipuram District. *Indian J Multidiscipl Dentistry* 2011;1(2): 115-118. 11
9. Bhat PK, Kadanakuppe S. Periodontal health status and oral hygiene practices of Iruliga tribal community residing at Ramanagar district, Karnataka, India. *J Int Oral Health* 2010;2(1):17-26.
10. Singh SK, Saha S, Jagannath GV, Singh P. Nature of Crime, Duration of Stay, Parafunctional Habits and Periodontal Status in Prisoners. *J Oral Health Comm Dent*. 2012;6(3):131-134.
11. Oral health surveys: basic methods. World Health Organization. 4th ed. Geneva: WHO; 1997.
12. Mullally BH. The Influence of Tobacco Smoking on the Onset of Periodontitis in Young Persons. *Tob Induc Dis* 2004;2(2):53-65.
13. Reddy S, Kaul S, Agrawal C, Prasad MG, Agnihotri J, Bhowmik N *et al*. Periodontal Status amongst Substance Abusers in Indian Population. *ISRN Dent* 2012;2012:460856.

14. Bouchard F. A health care needs assessment of federal inmates in Canada. *Can J Public Health*. 2004;95(Suppl):S9-S63.
15. Heidari E, Dickinson C, Wilson R, Fiske J. Oral health of remand prisoners in HMP Brixton, London. *Br Dent J* 2007;202(2):1-6.
16. Elvin-Lewis M. Plants used for teeth cleansing throughout the world. *J Prev Dent*. 1980; 6: 61-70.
17. Jain M, Mathur A, Sawla L, Choudhary G, Kabra K, Duraiswamy P. Oral health status of mentally disabled subjects in India. The greatest periodontal destruction was manifested in the older age groups. *J Oral Science* 2009;51(3): 333-340
18. Puscasu CG, Totolici I, Girdea M, Dumitriu AS. Carmen Hanganu. Tobacco Smoking and Periodontal Conditions in an Adult Population from Constanta, Romania. *OHDMBSC* 2009;8(3): 25-32.
19. Feldman RS, Bravacos JS, Rose CL. Association between smoking different tobacco products and periodontal disease indexes. *J Periodontol* 1983;54(8):481.
20. Awartani F, al-Jasser N. The effect of smoking on periodontal conditions assessed by CPITN. *Odontostomatol Trop* 1999;87:38-40.
21. Bali RK, Mathur VB, Talwar PP, Chanana HB. National Oral Health Survey and Fluoride Mapping 2002-2003 India. Dental Council of India and Ministry of Health and Family Welfare (Government of India), 2004.
22. Reddy V, Kondareddy CV, Siddanna S, Manjunath M. A survey on oral health status and treatment needs of life-imprisoned inmates in central jails of Karnataka, India. *Int Dent J* 2012;62(1):27-32.
23. Laxman VK, Annaji S. Tobacco use and its effects on the periodontium and periodontal therapy. *J Contemp Dent Pract* 2008;9(7):97-107.
24. Shimazaki Y, Saito T, Kiyohara Y, Kato I, Kubo M, Iida M *et al*. Relationship between drinking and periodontitis: the Hisayama study. *J Periodontol* 2005;76(9):1534-1541.
25. Picozzi A, Dworkin SF, Leeds JG, Nash J. Dental and associated attitudinal aspects of heroin addiction: a pilot study. *J Dent Res* 1972;51(3): 869.
26. Robinson PG, Acquah S, Gibson B. Drug users: oral health-related attitudes and behaviours. *Br Dent J* 2005;198(4):219-224.

Corresponding Address

Dr. Shailendra Singh Chaturvedi
 Reader
 Department of Periodontology,
 New Horizon Dental College and Research Institute,
 Bilaspur, Chhattisgarh, INDIA